

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/079193 A1

- (51) International Patent Classification⁷: **C07D 403/04**, A61K 31/506, A61P 35/00 (74) Agents: NACHSHEN, Neil, Jacob et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).
- (21) International Application Number: PCT/GB02/01445 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 26 March 2002 (26.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0107901.1 29 March 2001 (29.03.2001) GB (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): CYCLACEL LIMITED [GB/GB]; 12 St. James's Square, London SW1Y 4RB (GB).
- (72) Inventors; and (75) Inventors/Applicants (*for US only*): FISCHER, Peter, Martin [CH/GB]; Denley Lodge, 1 Arbirlot Road, Arbroath, Angus DD11 2EN (GB). WANG, Shudong [AU/GB]; Burnside Mill, Forfar, Angus, Scotland DD8 2RZ (GB). WOOD, Gavin [GB/GB]; 31 Millbank, Cupar, Fife, Scotland KY15 5DP (GB).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/079193 A1

(54) Title: INHIBITORS OF CYCLIN DEPENDENT KINASES AS ANTI-CANCER AGENT

(57) Abstract: The present invention relates to 2-substituted 4-heteroaryl-pyrimidines, their preparation, pharmaceutical compositions containing them and their use as inhibitors of cyclin-dependent kinases (CDKs) and hence their use in the treatment of proliferation disorders such as cancer, leukaemia, psoriasis and the like.

INHIBITORS OF CYCLIN DEPENDENT KINASES AS ANTI-CANCER AGENT

The present invention relates to 2-substituted 4-heteroaryl-pyrimidines, their preparation, pharmaceutical compositions containing them, and their use in the treatment of proliferative disorders such as cancer, leukemia, psoriasis and the like.

Introduction and Summary of the Prior art

Certain 4,5,6-Substituted-N-(substituted-phenyl)-2-pyrimidineamines having anti-asthmatic properties are disclosed in EP-A-233,461. Certain 4-heteroaryl-N-(3-substituted-phenyl)-2-pyridineamines possessing anti-proliferative properties and inhibiting protein kinases C, epidermal growth factor receptor-associated tyrosine protein kinase (EGF-R-TPK), as well as CDK1/cyclin B have been disclosed in WO95/09847 wherein the exemplified heteroaryl are pyridyl and indolyl.


J. Med. Chem. (1993) Vol. 36, pages 2716-2725, Paul, R. et al: discloses a further class of phenyl amino-pyrimidines possessing anti-inflammatory activity. These compounds include unsubstituted pyrrol groups, mono-substituted 2-thienyl groups and dimethyl-3-furyl groups at the 4-position of the pyrimidine ring.

It is an aim of the present invention to provide a further class of N-phenyl-2-pyrimidine anti-proliferative compounds. The compounds of the present invention have surprisingly been found to not to be inhibitors of protein kinase C. As discussed hereinafter, their activity may be demonstrated by inhibition of cell proliferation in cell lines and/or inhibition of cyclin dependent kinase enzymes.

Summary of the Invention

The first aspect of the present invention relates to compounds of general formula I:

5

**I**

wherein:

- 10 one of X^1 and X^2 is NR^{10} and the other of X^1 and X^2 is CR^9 ;

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, or CH=CH;

- 15 R^1 , R^2 , R^3 , R^9 and R^{10} are independently H, alkyl, aryl, aralkyl, heterocycle, halogeno, NO₂, CN, OH, alkoxy, aryloxy, $(R''')_nNH_2$, $(R''')_nNH-R'$, $(R''')_nN-(R')(R'')$, NH-aryl, N-(aryl)₂, COOH, COO-R', COO-aryl, CONH₂, CONH-R', CON-(R')(R''), CONH-aryl, CON-(aryl)₂, SO₃H, SO₂NH₂, CF₃, CO-R', or CO-aryl, wherein alkyl, aryl, aralkyl and heterocycle groups may be further substituted with one or more groups selected from halogeno, NO₂, CN, OH, O-methyl, NH₂, COOH, CONH₂ and
- 20 CF₃;

- R^4 , R^5 , R^6 , R^7 , and R^8 are independently from each other H, substituted or unsubstituted lower alkyl, halogeno, NO₂, CN, OH, substituted or unsubstituted alkoxy, NH₂, NH-R', N-(R')(R''), COOH, COO-R', CONH₂, CONH-R', CON-
- 25 (R')(R''), SO₃H, SO₂NH₂, or CF₃;

wherein R', R'' and R''' are each independently alkyl groups that may be the same or different and n is 0 or 1;

with the proviso that when R¹ and R² are H, X¹ is NH, X² is CH, and R³ is H, the

5 phenyl group is not

unsubstituted,

4-ethyl,

3-methyl,

3-(1,1,2,2-tetrafluoroethoxy),

10 3,4,5-trimethoxy,

when the other groups R⁴-R⁸ are H;

and pharmaceutically acceptable salts thereof.

Description of the Preferred Embodiments

15

As used herein the term "alkyl" includes both straight chain and branched alkyl groups having from 1 to 8 carbon atoms, e.g. methyl, ethyl propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl etc and the term "lower alkyl" is similarly used for groups having from 1 to 4 carbon atoms.

20

The term "aryl" is used to include groups having from 6 to 10 carbon atoms, e.g. phenyl, naphthyl etc.

The term "aralkyl" is used as a conjunction of the terms alkyl and aryl as given above.

25

Preferred compounds of formula I are those bearing a mono-, di- or trisubstituted pyrrol radical, attached to the pyrimidine ring through one of the ring carbon atoms. Preferably, the pyrrol radical is a pyrrol-3-yl group (i.e. X¹ is CR⁹ and X² is NR¹⁰, preferably NH) and is di- or tri-substituted.

The pyrrol group may be substituted by R^1 , R^2 , R^9 and R^{10} . Preferably, R^1 , R^2 and where appropriate R^9 and R^{10} are independently H, alkyl, aryl, aralkyl, heterocycle, halogeno, NO_2 , CN, OH, alkoxy, aryloxy, $(R''')_nNH_2$, $(R''')_nNH-R'$, $(R''')_nN-$
 5 $(R')(R'')$, NH-aryl, N-(aryl)₂, COOH, COO- R' , COO-aryl, CONH₂, CONH- R' , CON- $(R')(R'')$, CONH-aryl, CON-(aryl)₂, SO₃H, SO₂NH₂, CF₃, CO- R' , or CO-aryl, wherein alkyl, aryl, aralkyl and heterocycle groups may be further substituted with one or more groups selected from halogeno, NO_2 , CN, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃. Most preferably R^{10} is H.

10

More preferably, R^1 , R^2 and R^9 are each independently selected from H, halogeno, NO_2 , CN, $(R''')_nN-(R')(R'')$, CONH₂, a C₁₋₄ alkyl group and a heterocyclic group. Preferably, at least one, more preferably at least two or three of R^1 , R^2 and R^9 are not hydrogen.

15

Preferably, R^1 is H, CN, halogeno, nitro, alkylamino or a heterocyclic group. When R^1 is halogeno, it is preferably selected from chloro or bromo. When R^1 is alkylamino, it is preferably diethylaminomethyl or dimethylaminomethyl. When R^1 is a heterocyclic group it is preferably morpholin-4-ylmethyl or 4-methyl-piperazin-1-ylmethyl. Most preferably, R^1 is H or CN.
 20

Even more preferably, when R^1 is as preferably described, R^2 and R^9 are both lower alkyl, preferably methyl.

25 The group Z is preferably NH, NHSO₂ or NHCH₂, most preferably NH.

The phenyl substituents R^4 - R^8 are each independently selected from H, halogeno, nitro, amino, aminoalkyl, hydroxy, alkoxy, carbamoyl, sulfamyl, CN, N(R')(R''), C₁₋₄ alkyl and substituted C₁₋₄ alkyl.

More preferably, R^4 , R^5 , R^6 , R^7 , and R^8 are independently from each other H, unsubstituted lower alkyl, halogeno, NO_2 , CN, OH, N-(R')(R''), or CF_3 ; wherein R' , R'' and R''' are each independently alkyl groups that may be the same or different and n is 0 or 1;

Even more preferably, R^4 to R^8 are selected independently from H, F, NH_2 , NO_2 , OH, Cl, Br, I, CN, CH_2OH , CF_3 and dimethylamino. Within the preferences for R^4 to R^8 , R^4 and R^8 are most preferably hydrogen.

Thus, particularly preferred embodiments include 2-[N-(phenyl)]-4-(2,4-dimethylpyrrol-3-yl)pyrimidineamines in which the phenyl group is 2-, 3- or 4-substituted by at least one of H, F, NH_2 , NO_2 , OH, Cl, Br, I, CN, CH_2OH , CF_3 or OMe.

Within this particular embodiment, the phenyl group is preferably mono-substituted by F, NH_2 , NO_2 , OH, Cl, Br, I, CH_2OH , CN, CF_3 or OMe at any of the 2,3 or 4-positions, or di-substituted by 2,4-difluoro, 3,5-difluoro, 3,4-difluoro, 2,4-dichloro, 3,5-dichloro, 3,4-dichloro or 4-chloro-3-trifluoromethyl.

Further particularly preferred embodiments include 2-[N-(phenyl)]-4-(3,5-dimethyl-1H-pyrrole-2-carbonitrile)pyrimidineamines in which the phenyl group is 2-, 3- or 4-substituted by at least one of F, $NH(CH_3)_2$, NO_2 , OH, Cl, Br, I or CF_3 .

Within this particular embodiment, the phenyl group is preferably mono-substituted by F, $NH(CH_3)_2$, NO_2 , OH, I or CF_3 at any of the 3 or 4-positions, or di-substituted by 4-methyl-3-nitro, 3-iodo-4-methyl, 4-chloro-3-methyl, 3-hydroxy-4-methyl, 4-fluoro-3-methyl or 4-methyl-3-fluoro.

Further more particularly preferable embodiments include;

- 2-[N-(phenyl)]-4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is preferably mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position, preferably by a fluoro or NH(CH₃)₂ group.

5

- 2-[N-(phenyl)]-4-(2,4-dimethyl-5-halogeno-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is preferably mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 3 or 4-positions, preferably by a 4-fluoro or 3-nitro group, the halogeno group preferably being chloro or bromo.

10

- 2-[N-(phenyl)]-4-(2,4-dimethyl-5-dialkylaminoalkyl-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is preferably mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position, preferably by fluoro, the dialkylaminoalkyl group preferably being diethylaminomethyl or

15 dimethylaminomethyl.

- 2-[N-(phenyl)]-4-(2,4-dimethyl-5-(heterocycle)-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is preferably mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position, preferably by fluoro, the heterocycle group preferably being

20 5-morpholin-4-ylmethyl or 4-methyl-piperazin-1-ylmethyl.

Most preferably, the compounds of the present invention are selected from;

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

25 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine

(3,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine

(4-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine

(3,5-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine

4-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol

- 3-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
 (2,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (2,4-Dichloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (4-Chloro-3-trifluoromethyl-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 5 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-trifluoromethyl-phenyl)-amine
 (3-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 N-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-
 10 diamine
 (3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-fluoro-4-iodo-phenyl)-amine
 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 15 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 20 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 25 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 5 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- [4-(3,5-Dimethyl-1H-pyrrol-2-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 10 (4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine
- [4-(5-Amino-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 15 [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 20 [4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- {4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine
- 25

The structures of the above-mentioned compounds are illustrated in Figure 1.

Particularly preferred compounds observed are those to be CDK inhibitors having IC₅₀ for cdk2/cyclinE of less than 5μM(±0.05), including;

- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine
 (3,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 5 (4-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (3,5-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 4-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
 3-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
 10 (3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-fluoro-4-iodo-phenyl)-amine
 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 15 carbonitrile
 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-
 carbonitrile
 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 20 carbonitrile
 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-
 carbonitrile
 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 25 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile

4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

5 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide

(4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine

10 [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

15 [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
[4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

20 [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine, and

{4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine.

25 Of these compounds, more preferred are those to be CDK inhibitors having IC_{50} for cdk2/cyclinE of less than $1\mu M(\pm 0.05)$, including;

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine
- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
- 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 5 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 10 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 15 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 20 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 25 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

5 [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine, and

10 [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine.

Of these, even more preferred are compounds are those having IC_{50} for cdk2/cyclinE of less than $0.5\mu M$ (± 0.05), being;

15

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine

3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

20 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

25 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

5 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide

[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine, and

10 [4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine.

The following compounds are observed to be particularly effective anti-proliferative agents, as demonstrated by cell-based assays:

15 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine

(3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine

3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

20 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

25 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

- 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 5 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 10 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- 15 (4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- 20 [4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine, and
- {4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine.
- 25

The compounds of formula I have been found to possess anti-proliferative activity and are therefore believed to be of use in the treatment of proliferative disorders such as cancers, leukaemias and other disorders associated with uncontrolled cellular

proliferation such as psoriasis and restenosis. As defined herein, an anti-proliferative effect within the scope of the present invention may be demonstrated by the ability to inhibit cell proliferation in an *in vitro* whole cell assay, for example using any of the cell lines A549, HT29, Saos-2, HeLa or MCF-7, or by showing inhibition of a CDK enzyme (such as CDK2 or CDK4) in an appropriate assay. These assays, including methods for their performance, are described in more detail in Example 3. Using such cell line and enzymes assays it may be determined whether a compound is anti-proliferative in the context of the present invention.

Without wishing to be bound by theory, the compounds of the present invention are believed to exert their anti-proliferative effect in a non-protein kinase C (PKC) dependent manner. Many of the compounds inhibit cyclin-dependent kinase enzymes (CDKs) that have been shown to be involved in cell cycle control. These CDKs include CDK2 and CDK4 and particularly their respective interactions with cyclin E and cyclin D1. These compounds of the present invention are further believed to be advantageous in being selective for CDK enzymes implicated in proliferative diseases. By the term "selective" it is meant that although possible having some inhibitory effect on another enzyme (such as PKC), the compound is preferentially effective against an enzyme implicated in proliferative diseases.

20


The compounds of the invention may inhibit any of the steps or stages in the cell cycle, for example, formation of the nuclear envelope, exit from the quiescent phase of the cell cycle (G0), G1 progression, chromosome decondensation, nuclear envelope breakdown, START, initiation of DNA replication, progression of DNA replication, termination of DNA replication, centrosome duplication, G2 progression, activation of mitotic or meiotic functions, chromosome condensation, centrosome separation, microtubule nucleation, spindle formation and function, interactions with microtubule motor proteins, chromatid separation and segregation, inactivation of mitotic functions, formation of contractile ring, and cytokinesis functions. In particular, the

compounds of the invention may influence certain gene functions such as chromatin binding, formation of replication complexes, replication licensing, phosphorylation or other secondary modification activity, proteolytic degradation, microtubule binding, actin binding, septin binding, microtubule organising centre nucleation activity and
5 binding to components of cell cycle signalling pathways.

A further embodiment of the present invention therefore relates to the use of one or more compounds of formula I in the treatment of proliferative disorders. Preferably, the proliferative disorder is a cancer or leukaemia. The term proliferative disorder is
10 used herein in a broad sense to include any disorder that requires control of the cell cycle, for example cardiovascular disorders such as restenosis and cardiomyopathy, auto-immune disorders such as glomerulonephritis and rheumatoid arthritis, dermatological disorders such as psoriasis, anti-inflammatory, anti-fungal, antiparasitic disorders such as malaria, emphysema and alopecia. In these disorders,
15 the compounds of the present invention may induce apoptosis or maintain stasis within the desired cells as required.

In a particularly preferred embodiment, the invention relates to the use of one or more compounds of formula I in the treatment of a CDK dependent or sensitive disorder.
20 CDK dependent disorders are associated with an above normal level of activity of one or more CDK enzymes. Such disorders preferably associated with an abnormal level of activity of CDK2 and/or CDK4. A CDK sensitive disorder is a disorder in which an aberration in the CDK level is not the primary cause, but is downstream of the primary metabolic aberration. In such scenarios, CDK2 and/or CDK4 can be said to
25 be part of the sensitive metabolic pathway and CDK inhibitors may therefore be active in treating such disorders. Such disorders are preferably cancer or leukaemic disorders.

A second aspect of the present invention relates to the use of a compound of formula



II

5 wherein:

one of X^1 and X^2 is NH and the other of X^1 and X^2 is CR^9 ;

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, or CH=CH;

- 10 R^1 , R^2 , R^3 and R^9 are independently H, alkyl, aryl, aralkyl, heterocycle, halogeno, NO₂, CN, OH, alkoxy, aryloxy, NH₂, NH-R', N-(R')(R''), NH-aryl, N-(aryl)₂, COOH, COO-R', COO-aryl, CONH₂, CONH-R', CON-(R')(R''), CONH-aryl, CON-(aryl)₂, SO₃H, SO₂NH₂, CF₃, CO-R', or CO-aryl, wherein alkyl, aryl, aralkyl and heterocycle groups may be further substituted with one or more groups selected from halogeno,
- 15 NO₂, CN, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

- R^4 , R^5 , R^6 , R^7 , and R^8 are independently from each other H, substituted or unsubstituted lower alkyl, halogeno, NO₂, CN, OH, substituted or unsubstituted alkoxy, NH₂, NH-R', N-(R')(R''), COOH, COO-R', CONH₂, CONH-R', CON-
- 20 (R')(R''), SO₃H, SO₂NH₂, or CF₃;

wherein R' and R'' are each independently alkyl groups that may be the same or different;

with the proviso that when R^1 and R^2 are H, X^1 is NH, X^2 is CH, and R^3 is H, the phenyl group is not

3-(1,1,2,2- tetrafluoroethoxy), or

3,4,5-trimethoxy,

5 when the other groups R^4 - R^8 are H;

and pharmaceutically acceptable salts thereof;

in the manufacture of a medicament for use in the treatment of a proliferative disease.

10 The term "proliferative disorder" has been previously discussed and the same definition applies to the second aspect of the invention.

A further aspect of the present invention relates to the use of the compounds of formula II in the manufacture of a medicament for use in the treatment of antiviral infections. Such viral infections include VZV, HSV type 1 and 2 and HIV.
15 Preferably, the compounds are of use in the treatment of HIV and HIV related disorders.

The preferred embodiments of these further aspects of the invention are identical to those described above in respect of the first aspect.

20

In a particularly preferred embodiment, the one or more compounds of the invention are administered in combination with one or more other anticancer agents. In such cases, the compounds of the invention may be administered consecutively, simultaneously or sequentially with the one or more other anticancer agents.

25

As used herein the phrase "manufacture of a medicament" includes the use of a compound of formula I directly as the medicament in addition to its use in a screening programme for further anti-proliferative agents or in any stage of the manufacture of such a medicament.

The compounds of the present invention (first and seconds aspects) can be present as salts or esters, in particular pharmaceutically acceptable salts or esters.

5 Pharmaceutically acceptable salts of the compounds of the invention (first and seconds aspects) include suitable acid addition or base salts thereof. A review of suitable pharmaceutical salts may be found in Berge et al, J Pharm Sci, 66, 1-19 (1977). Salts are formed, for example with strong inorganic acids such as mineral acids, e.g. sulphuric acid, phosphoric acid or hydrohalic acids; with strong organic
10 carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted (e.g., by halogen), such as acetic acid; with saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid; with aminoacids, for example aspartic or
15 glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted (for example, by a halogen) such as methane- or p-toluene sulfonic acid.

Esters are formed either using organic acids or alcohols/hydroxides, depending on the
20 functional group being esterified. Organic acids include carboxylic acids, such as alkanecarboxylic acids of 1 to 12 carbon atoms which are unsubstituted or substituted (e.g., by halogen), such as acetic acid; with saturated or unsaturated dicarboxylic acid, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric
25 acid; with aminoacids, for example aspartic or glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted (for example, by a halogen) such as methane- or p-toluene sulfonic acid. Suitable hydroxides include inorganic hydroxides, such as sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminium hydroxide. Alcohols

include alkanealcohols of 1-12 carbon atoms which may be unsubstituted or substituted, e.g. by a halogen).

In all aspects of the present invention previously discussed, the invention includes,
5 where appropriate all enantiomers and tautomers of compounds of formula I. The man skilled in the art will recognise compounds that possess an optical properties (one or more chiral carbon atoms) or tautomeric characteristics. The corresponding enantiomers and/or tautomers may be isolated/prepared by methods known in the art.

10 The invention furthermore relates to the compounds of or of use in the present invention (first and seconds aspects) in their various crystalline forms, polymorphic forms and (an)hydrous forms. It is well established within the pharmaceutical industry that chemical compounds may be isolated in any of such forms by slightly varying the method of purification and or isolation from the solvents used in the
15 synthetic preparation of such compounds.

The invention further includes the compounds (first and seconds aspects) of or of use in the present invention in prodrug form. Such prodrugs are generally compounds of formula I wherein one or more appropriate groups have been modified such that the
20 modification may be reversed upon administration to a human or mammalian subject. Such reversion is usually performed by an enzyme naturally present in such subject, though it is possible for a second agent to be administered together with such a prodrug in order to perform the reversion in vivo. Examples of such modifications include ester (for example, any of those described above), wherein the reversion may
25 be carried out by an esterase etc. Other such systems will be well known to those skilled in the art.

The present invention also encompasses pharmaceutical compositions comprising the compounds of the invention (first and seconds aspects). In this regard, and in

particular for human therapy, even though the compounds of the present invention (including their pharmaceutically acceptable salts, esters and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutical carrier, excipient or diluent selected with regard to the intended route of administration and standard pharmaceutical practice.

Thus, the present invention also relates to pharmaceutical compositions comprising one or more compounds of formula I or II or pharmaceutically acceptable salts or esters thereof, together with at least one pharmaceutically acceptable excipient, diluent or carrier.

By way of example, in the pharmaceutical compositions of the present invention, the compounds of the invention may be admixed with any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), and/or solubilising agent(s). Examples of such suitable excipients for the various different forms of pharmaceutical compositions described herein may be found in the "Handbook of Pharmaceutical Excipients, 2nd Edition, (1994), Edited by A Wade and PJ Weller.

The pharmaceutical compositions of the present invention may be adapted for oral, rectal, vaginal, parenteral, intramuscular, intraperitoneal, intraarterial, intrathecal, intrabronchial, subcutaneous, intradermal, intravenous, nasal, buccal or sublingual routes of administration.

For oral administration, particular use is made of compressed tablets, pills, tablets, gellules, drops, and capsules. Preferably, these compositions contain from 1 to 250 mg and more preferably from 10-100 mg, of active ingredient per dose.

Other forms of administration comprise solutions or emulsions which may be injected intravenously, intraarterially, intrathecally, subcutaneously, intradermally,

intraperitoneally or intramuscularly, and which are prepared from sterile or sterilisable solutions. The pharmaceutical compositions of the present invention may also be in form of suppositories, pessaries, suspensions, emulsions, lotions, ointments, creams, gels, sprays, solutions or dusting powders.

5

An alternative means of transdermal administration is by use of a skin patch. For example, the active ingredient can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin. The active ingredient can also be incorporated, at a concentration of between 1 and 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be required.

10

Injectable forms may contain between 10 - 1000 mg, preferably between 10 - 250 mg, of active ingredient per dose.

15

Compositions may be formulated in unit dosage form, i.e., in the form of discrete portions containing a unit dose, or a multiple or sub-unit of a unit dose.

20

A person of ordinary skill in the art can easily determine an appropriate dose of one of the instant compositions to administer to a subject without undue experimentation. Typically, a physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The dosages disclosed herein are exemplary of the average case. There can of course be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

25

In an exemplary embodiment, one or more doses of 10 to 150 mg/day will be administered to the patient for the treatment of malignancy.

The pharmaceutical compositions of the invention may further comprise one or more additional anticancer agents, for example, existing anticancer drugs available on the market.


- 5 Anticancer drugs in general are more effective when used in combination. In particular, combination therapy is desirable in order to avoid an overlap of major toxicities, mechanism of action and resistance mechanism(s). Furthermore, it is also desirable to administer most drugs at their maximum tolerated doses with minimum time intervals between such doses. The major advantages of combining
10 chemotherapeutic drugs are that it may promote additive or possible synergistic effects through biochemical interactions and also may decrease the emergence of resistance in early tumor cells which would have been otherwise responsive to initial chemotherapy with a single agent. An example of the use of biochemical interactions in selecting drug combinations is demonstrated by the administration of leucovorin to increase the
15 binding of an active intracellular metabolite of 5-fluorouracil to its target, thymidylate synthase, thus increasing its cytotoxic effects.

Numerous combinations are used in current treatments of cancer and leukemia. A more extensive review of medical practices may be found in "Oncologic Therapies"
20 edited by E. E. Vokes and H. M. Golomb, published by Springer.

Beneficial combinations may be suggested by studying the growth inhibitory activity of the test compounds with agents known or suspected of being valuable in the treatment of a particular cancer initially or cell lines derived from that cancer. This
25 procedure can also be used to determine the order of administration of the agents, i.e. before, simultaneously, or after delivery. Such scheduling may be a feature of all the cycle acting agents identified herein.

The compounds of this invention (I) can be synthesised, for example, by an adaptation of the Traube synthesis (A.R. Katritzky, I. Taher, *Can. J. Chem.* 1986, 64, 2087 and references cited therein), *i.e.* by condensation between 1,3-dicarbonyl compounds 1 or acrylates 2 or 3, and amidine 4, as shown in *Scheme 1*.


5



Scheme 1


- 10 The dicarbonyl compounds I in turn can be prepared by many methods known in the art (J. March, *In: Advanced Organic Chemistry: Reactions, Mechanism, and Structure*, 4th Ed., John Wiley & Sons, Inc., New York, 1992, p. 1283). Acrylates 2 and 3, which are particularly suitable for the purposes of this invention, are obtained from

heterocyclic methyl ketones **5** by condensation with *tert*-butoxybis(dimethylamino)methane **6** (Scheme 2).



Scheme 2


- 5 The diamino compounds **4** will be amidines **4a** or guanidines **4b**, depending on the definition of Z in general structure I. Amidines (HN=CRNH₂) can be obtained from readily available amine precursors by condensation with *e.g.* ketenimines, or by addition of ammonia to suitable nitriles or imidates. Guanidines **4b** (Scheme 3) can be elaborated by a number of methods known in the art. For the purposes of this
- 10 invention, the most useful route is amination of cyanamide **8** with anilines **9**.



Scheme 3

In the case where 5 is a pyrrole, two systems can apply (refer *Scheme 4*), i.e. the acetyl group which is used to generate the pyrimidine precursors 2 and 3 is either in the pyrrole 3-position (5: $X^1 = CR^9$, $X^2 = NH$; i.e. structure 5b) or in the pyrrole 2-position ($X^1 = NH$, $X^2 = CR^9$; i.e. structure 5c).

5



Scheme 4'

In both cases the pyrrole rings can be assembled using methods known in the art. Particularly relevant is a modification of the Knorr synthesis (refer, e.g. J. A. Joule, G. F. Smith, *Heterocyclic Chemistry*, 2nd Ed., Van Nostrand Reinhold (UK) Co. Ltd., 1978, pp.213-215). For the pyrrol-3-yl system, activated (i.e. $R^1 = COOEt$, CN, etc.) carbonyl compounds **10** are first nitrosylated. The resulting oximes **11** are condensed with dicarbonyl compounds **12** in the presence of e.g. zinc-acetic acid or aqueous dithionite, with formation of the reactive α -aminocarbonyl intermediate **13**. The R^1 substituent (e.g. $COOEt$, CN) in the resulting 3-acetylpyrroles **5b** can be further manipulated, either directly, or in the context of intermediates 2 or 3, or in the pyrrolopyrimidine products I. Thus decarboxylation ($R^1 = COOEt$) will give products with $R^1 = H$, oxidation ($R^1 = CN$) will afford products with $R^1 = CONH_2$, etc.

Furthermore, products with $R^1 = H$ can be transformed into various derivatives, particularly through electrophilic substitution. Thus derivatives where R^1 is, for example, a halogen, nitro, amino, alkyl, alkylamino, *etc.*, group can be obtained readily. In the case of the pyrrol-2-yl system an analogous situation arises, here an
5 activating group needs to be present in the carbonyl component 15 (*e.g.* $R^9 = COOEt$, CN , *etc.*). This is condensed with oximes 16 (derived from dicarbonyl compounds 14), again with formation of the intermediate 17. The R^9 substituent in products 5c or derivatives can be manipulated in the same way as the R^1 group in the pyrrol-3-yl system discussed above.

10

Alternatively, compounds of general structure I can be obtained from suitable pyrimidine precursors directly, *e.g.* from 2,4-disubstituted (halogen, amine, *etc.*) pyrimidines by successive substitution reactions.

15 The present invention will now be described by way of example and with reference to the following figure:

Figure 1 shows the chemical structure of compounds according to the invention.

20 Examples

Abbreviations

LC-MS, liquid chromatography-mass spectrometry; NMR, nuclear magnetic resonance spectroscopy; r.t. room temperature; PE, petroleum ether (40-60 °C boiling
25 fraction); DMSO, dimethylsulfoxide.


General

NMR spectra were recorded using a Bruker DPX-300 instrument. Chemical shifts are reported in ppm (δ) from tetramethylsilane. EM Kieselgel 60 (0.040-0.063 mm) was

used for flash column chromatography. Melting points were determined with a LEICA testo-720 electrothermometer and are uncorrected. Compound numbers are shown in brackets, where appropriate.

5 Example 1

3-Dimethylamino-1-(2,4-dimethyl-1H-pyrrol-3-yl)-propenone



A mixture of 1-(2,4-dimethyl-1H-pyrrol-3-yl)-ethanone (2 g, 15 mmol) in 5 mL of 1,1-bis-dimethylamino-3,3-dimethyl-butan-2-one was heated at 100 °C for 22 h. The precipitates of the reaction mixture were slurried in EtOAc/PE with chilling. The crude product was filtered, washed with EtOAc/PE, and dried *in vacuo* to afford the title compound as a purple solid (2.6 g). ¹H-NMR (CDCl₃) δ: 2.25 (s, 6H, CH₃), 2.45 (s, 6H, CH₃), 5.46 (d, 1H, *J* = 12.6 Hz, CH), 6.35 (s, 1H, pyrrolyl-H), 7.63 (d, 1H, *J* = 12.6 Hz, CH).

15

Example 2

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [1]

To a mixture of 3-dimethylamino-1-(2,4-dimethyl-1H-pyrrol-3-yl)-propenone (1 mmol, 0.19 g) and 4-fluorophenyl guanidine nitrate (2 mmol, 0.44 g) in 2-methoxyethanol (5 mL) was added NaOH (40 mg). The reaction mixture was heated at 100-120 °C under N₂ for 6 h. The solvent was evaporated to dryness and the residue was purified by flash chromatography (1:2 EtOAc/PE). Recrystallisation from EtOAc/PE afforded the title compound (174 mg, 62 %) as brown crystals. ¹H-NMR (CDCl₃) δ: 2.21 (s, 3H, CH₃), 2.43 (s, 3H, CH₃), 6.33 (s, 1H, pyrrolyl-H), 6.73 (d, 1H, *J* = 5.3 Hz, pyrimidinyl-H), 7.00 (m, 2H, Ph-H), 7.79 (m, 2H, Ph-H), 8.28 (d, 1H, *J* = 5.3 Hz, pyrimidinyl-H), 9.16 (s, 1H, NH), 10.59 (s, 1H, NH).

25

The following compounds were prepared in a manner analogous to that described above:

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine [2]

- 5 Yellow-orange solid. M.p. 197-199 °C. LC-MS: m/z = 310 (M+1). $C_{16}H_{15}N_5O_2$ requires C, 62.12; H, 4.89; N, 22.64; found C, 62.61; H, 4.99; N, 22.20. 1H -NMR ($CDCl_3$) δ : 2.71 (d, 6H, CH_3), 7.05(d, 1H, J = 5.3 Hz, pyrimidinyl-H), 7.47 (m, 2H, Ph-H), 7.78 (m, 1H, Ph-H), 7.81 (s, 1H, Ar-H), 8.07 (m, 1H, Ph-H), 8.51 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 8.99 (br. s, 1H, NH), 9.91 (br. s, 1H, NH).

10

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine [3]

1H -NMR ($CDCl_3$) δ : 2.26 (s, 3H, CH_3), 2.48 (s, 3H, CH_3), 6.80 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 7.47 (m, 2H, Ph-H), 7.57 (m, 2H, Ph-H), 7.23 (s, 1H, pyrrolyl-H), 8.32 (d, 1H, J = 5.3 Hz, pyrimidinyl-H).

15

(3,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [4]

1H -NMR ($CDCl_3$) δ : 2.23 (s, 3H, CH_3), 2.46 (s, 3H, CH_3), 6.42 (m, 1H, pyrrolyl-H), 6.77 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 7.11 (m, 1H, Ph-H), 7.41 (m, 1H, Ph-H), 8.05 (m, 1H, Ph-H), 8.29 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 9.21 (s, 1H, Ph-H), 10.46 (br. s, 1H, NH).

20

(4-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [5]

M.p. 219-223 °C. MS: $[M+H]^+$ = 299.4 ($C_{16}H_{15}ClN_4$ requires 298.8). 1H -NMR ($DMSO-d_6$) δ : 2.19 (s, 3H, CH_3), 2.42 (s, 3H, CH_3), 6.48 (s, 1H, pyrrolyl-H), 6.82 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 7.31 (d, 2H, J = 8.7Hz, Ph-H), 7.84 (d, 2H, J = 8.7Hz, Ph-H), 8.34 (d, 1H, J = 5.3 Hz, pyrimidinyl-H), 9.45 (s, 1H, Ph-H), 10.72 (br. s, 1H, NH).

25

(3,5-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [6]

M.p. 153.3-156.8 °C. MS: $[M+H]^+ = 303.6$ ($C_{16}H_{14}F_2N_4$ requires 300.3). 1H -NMR (CD_3OD) δ : 2.25 (s, 3H, CH_3), 2.47 (s, 3H, CH_3), 6.42-6.48 (m, 2H, pyrrolyl-H and Ph-H), 6.80 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.44-7.48 (m, 2H, Ph-H), 8.31 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H).

5

4-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol [7]

M.p. 189.5-193.4 °C. MS: $[M+H]^+ = 281.9$ ($C_{16}H_{16}N_4O$ requires 280.3). 1H -NMR (CD_3OD) δ : 2.23 (s, 3H, CH_3), 2.43 (s, 3H, CH_3), 6.44 (s, 1H, pyrrolyl-H), 6.75-6.78 (m, 3H, pyrimidinyl-H and Ph-H), 7.39 (d, 2H, $J = 8.8$ Hz, Ph-H), 8.17 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H).

10

3-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol [8]

M.p. 169.0-174.6 °C. MS: $[M+H]^+ = 281.3$ ($C_{16}H_{16}N_4O$ requires 280.3). 1H -NMR (CD_3OD) δ : 2.26 (s, 3H, CH_3), 2.47 (s, 3H, CH_3), 6.44-6.48 (m, 2H, pyrrolyl-H, Ph-H), 6.84 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.05-7.10 (m, 2H, Ph-H), 7.32 (m, 1H, Ph-H), 8.25 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H).

15

(2,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [9]

M.p. 219-220 °C. MS: $[M+H]^+ = 302.6$ ($C_{16}H_{14}F_2N_2$ requires 300.3). 1H -NMR ($DMSO-d_6$) δ : 2.10 (s, 3H, CH_3), 2.36 (s, 3H, CH_3), 6.43 (s, 1H, pyrrolyl-H), 6.77 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.06 (m, 1H, Ph-H), 7.27 (m, 1H, Ph-H), 7.66 (m, 1H, Ph-H), 8.25 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 8.71 (s, 1H, Ph-H), 10.70 (br. s, 1H, NH).

20

(2,4-Dichloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [10]

M.p. 158.5-159.7 °C. MS: $[M+H]^+ = 335.4$ ($C_{16}H_{14}Cl_2N_4$ requires 333.2). 1H -NMR ($DMSO-d_6$) δ : 2.18 (s, 3H, CH_3), 2.38 (s, 3H, CH_3), 6.51 (s, 1H, pyrrolyl-H), 6.90 (d, 1H, $J = 5.5$ Hz, pyrimidinyl-H), 7.46 (m, 1H, Ph-H), 7.71 (m, 1H, Ph-H), 8.05 (m, 1H,

25

Ph-H), 8.36 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 8.49 (s, 1H, Ph-H), 10.80 (br. s, 1H, NH).

5 *(4-Chloro-3-trifluoromethyl-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [11]*

M.p. 187.7-190.7 °C. MS: $[M+H]^+ = 368.6$ ($C_{17}H_{14}ClF_3N_4$ requires 366.8). 1H -NMR (DMSO- d_6) δ : 2.19 (s, 3H, CH_3), 2.42 (s, 3H, CH_3), 6.50 (s, 1H, pyrrolyl-H), 6.89 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.61 (m, 1H, Ph-H), 8.08 (m, 1H, Ph-H), 8.39-8.42 (m, 2H, Ph-H and pyrimidinyl-H), 9.79 (s, 1H), 10.80 (br. s, 1H, NH).

10

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine [12]

M.p. 165.6-167.9 °C. MS: $[M+H]^+ = 332.9$ ($C_{17}H_{15}F_3N_4$ requires 332.3). 1H -NMR (DMSO- d_6) δ : 2.26 (s, 3H, CH_3), 2.49 (s, 3H, CH_3), 6.56 (s, 1H, pyrrolyl-H), 6.94 (d, 15 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.67 (d, 2H, $J = 8.5$ Hz, Ph-H), 8.09 (d, 2H, $J = 8.5$ Hz, Ph-H), 8.45 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 9.82 (s, 1H, NH).

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-trifluoromethyl-phenyl)-amine [13]

20 M.p. 152.7-154.3 °C. MS: $[M+H]^+ = 332.6$ ($C_{17}H_{15}F_3N_4$ requires 332.3). 1H -NMR (DMSO- d_6) δ : 2.26 (s, 3H, CH_3), 2.48 (s, 3H, CH_3), 6.56 (s, 1H, pyrrolyl-H), 6.92 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 7.29 (m, 1H, Ph-H), 7.55 (m, 1H, Ph-H), 8.03 (m, 1H, Ph-H), 8.42-7.45 (m, 2H, pyrimidinyl-H and Ph-H), 9.73 (s, 1H, NH), 10.83 (br. s, 1H, NH).

25

(3-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [14]

M.p. 140.4-144.2 °C. MS: $[M+H]^+ = 299.5$ ($C_{16}H_{15}ClN_4$ requires 298.8). 1H -NMR (DMSO- d_6) δ : 2.21 (s, 3H, CH_3), 2.44 (s, 3H, CH_3), 6.51 (s, 1H, pyrrolyl-H), 6.85 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 6.94 (d, 1H, $J = 7.6$ Hz, Ph-H), 7.28 (t, 1H, $J = 8.1$ Hz,

Ph-H), 7.61 (d, 1H, $J = 8.2$ Hz, Ph-H), 8.19 (s, 1H, Ph-H), 8.37 (d, 1H, $J = 5.3$ Hz, pyrimidinyl-H), 9.55 (s, 1H, NH), 10.78 (br. s, 1H, NH).

5 *N*-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-*N*',*N*'-dimethyl-benzene-1,4-diamine [15]


M.p. 179.9-182.1 °C. MS: $[M+H]^+ = 307.3$ ($C_{18}H_{21}N_5$ requires 307.4). 1H -NMR ($CDCl_3$) δ : 2.25 (s, 3H, CH_3), 2.46 (s, 3H, CH_3), 2.91 (s, 6H, CH_3), 6.46 (s, 1H, pyrrolyl-H), 6.70 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 6.78 (dd, 2H, $J = 6.8, 2.2$ Hz, Ph-H), 6.79 (br. s, 1H, NH), 7.45 (dd, 2H, $J = 6.8, 2.2$ Hz, Ph-H), 7.80 (br. s, 1H, NH), 8.28 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H).

(3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine [16]

M.p. 185.0-187.4 °C. MS: $[M+H]^+ = 423.9$ ($C_{16}H_{14}ClIN_4$ requires 424.7). 1H -NMR ($DMSO-d_6$) δ : 2.18 (s, 3H, CH_3), 2.41 (s, 3H, CH_3), 6.48 (s, 1H, pyrrolyl-H), 6.84 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.40 (dd, 1H, $J = 8.8, 2.4$ Hz, Ph-H), 7.75 (d, 1H, $J = 8.8$ Hz, Ph-H), 8.34 (m, 1H, Ph-H), 8.36 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.61 (s, 1H, NH), 10.75 (s, 1H, NH).

20 *[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-fluoro-4-iodo-phenyl)-amine* [17]

M.p. 200-202 °C. MS: $[M+H]^+ = 407.4$ ($C_{16}H_{14}FIN_4$ requires 408.2). 1H -NMR ($DMSO-d_6$) δ : 2.18 (s, 3H, CH_3), 2.40 (s, 3H, CH_3), 6.48 (s, 1H, pyrrolyl-H), 6.84 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.35 (m, 1H, Ph-H), 7.64 (t, 1H, $J = 8.0$ Hz, Ph-H), 8.02 (dd, 1H, $J = 12.0, 2.2$ Hz, Ph-H), 8.36 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.65 (s, 1H, NH), 10.75 (s, 1H, NH).

Example 3***4-(3-Dimethylamino-acryloyl)-3,5-dimethyl-1H-pyrrole-2-carbonitrile***

Ethyl cyanoacetate (10 mL, 94 mmol) was diluted with AcOH (20 mL) and the solution was cooled to -10°C (ice-MeOH bath). NaNO_2 (6.5 g, 94 mmol) was dissolved in H_2O (10 mL) and the solution was added dropwise over a period of 40 min, keeping the internal temperature $< 0^{\circ}\text{C}$. After completion of the addition, the reaction mixture was stirred for 1 h with cooling. It was then warmed to room temperature and stirred for a further 3 h. The mixture was diluted with acetic acid (50 mL) and H_2O (50 mL). Pentane-2,4-dione (10.6 mL, 103 mmol) was added and the mixture heated to $\sim 75^{\circ}\text{C}$. To this reaction mixture Zn powder (6.9 g, 105 mmol) was added in portions over a period of 30 min at such a rate as to maintain the internal temperature $< 90^{\circ}\text{C}$. The reaction mixture was then heated for a further 30 min before pouring into H_2O (1 mL). From the reaction mixture 3,5-dimethyl-1H-pyrrole-2-carbonitrile (3.67 g) was filtered as an off-white solid. The filtrate was extracted with EtOAc (3×500 mL). The combined organic extracts were washed (brine) and dried (MgSO_4). The solvent was evaporated to a brown oil, which was purified by chromatography (100 g SiO_2 , eluted with 4:1 heptane / EtOAc) to afford a further crop (4.41 g) of this product as a pale yellow solid (total yield 72 %).

3,5-Dimethyl-1H-pyrrole-2-carbonitrile (1.2 g, 10 mmol) was dissolved in anhydrous 1,2-dichloroethane (15 mL) and AlCl_3 (2.93 g, 22 mmol) was added proportion-wise. The reaction vessel was purged with N_2 and was cooled in an ice-water bath. AcCl (0.71 mL, 10 mmol) was added dropwise and the mixture was stirred for 1 h with cooling and for a further 3 h at room temperature. The reaction mixture was quenched by careful addition of 2 M aq HCl . The acidity of the mixture was adjusted to approximately pH 6 by addition of NaHCO_3 . After separation of the organic phase, the aqueous phase was extracted with EtOAc ($3 \times 100\text{mL}$). The combined organic phases were washed (H_2O , then brine), dried (MgSO_4), and filtered.

The solvent was evaporated to afford of 4-acetyl-3,5-dimethyl-1*H*-pyrrole-2-carbonitrile (1.42 g, 88 %) as a pale tan solid. ¹H-NMR (CDCl₃) δ: 2.44 (s, 3H, CH₃), 2.45 (s, 3H, CH₃), 2.54 (s, 3H, CH₃), 8.75 (br. s, 1H, NH).

4-Acetyl-3,5-dimethyl-1*H*-pyrrole-2-carbonitrile (1.38 g, 8.51 mmol) was
5 suspended in 1,1-bis-dimethylamino-3,3-dimethyl-butan-2-one (1.3 mL) and heated at
75 °C for 42 h. The reaction mixture was evaporated to dryness and the residue was
purified by SiO₂ chromatography (heptane / EtOAc) to afford the title compound (1.2
g, 65 %) as a pale tan solid. ¹H-NMR (DMSO-d₆) δ: 2.21 (s, 3H, CH₃), 2.31 (s, 3H,
CH₃), 3.32 (s, 6H, CH₃), 5.22 (d, 1H, *J* = 12.4 Hz, CH), 7.47 (d 1H, *J* = 12.4 Hz; CH),
10 11.96 (br. s, 1H, NH).

Example 4

3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile [31]

15 To a mixture of 4-(3-dimethylamino-acryloyl)-3,5-dimethyl-1*H*-pyrrole-2-carbonitrile
(1.0 mmol, 0.22 g) and 3-nitrophenyl guanidine nitrate (1.5 mmol, 0.36 g) in 2-
methoxyethanol (5 mL) was added K₂CO₃ (138 mg, 1.0 mmol). The reaction mixture
was heated at 120 °C under N₂ for 18 h. The solvent was evaporated to dryness and
the residue was purified by flash chromatography (1:2 EtOAc / heptane) to afford the
20 title compound as a light-yellow solid. M.p. 258-259 °C. MS: [M+H]⁺ = 336.1
(C₁₇H₁₄N₆O₂ requires 334.3). ¹H-NMR (CD₃OD) δ: 2.39 (s, 3H, CH₃), 2.49 (s, 3H,
CH₃), 6.94 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 7.50 (t, 1H, *J* = 8.3 Hz, Ph-H), 7.81
(m, 1H, Ph-H), 7.94 (m, 1H, Ph-H), 8.45 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 8.94 (t,
1H, *J* = 2.2 Hz, Ph-H).

25

The following compounds were prepared in a manner analogous to that described
above:

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
[32]

MS: $[M+H]^+ = 307.7$ ($C_{17}H_{14}FN_5$ requires 307.3). 1H -NMR (DMSO- d_6) δ : 2.30 (s, 3H, CH_3), 2.40 (s, 3H, CH_3), 6.84 (d, 1H, $J = 5.0$ Hz, pyrimidinyl -H), 7.00 (m, 2H, Ph-H), 7.73 (m, 2H, Ph-H), 8.40 (d, 1H, $J = 5.5$ Hz, pyrimidinyl -H), 9.46 (s, 1H, NH), 12.19 (br. s, 1H, NH).

4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [33]

M.p. 272-276 °C. MS: $[M+H]^+ = 305.8$ ($C_{17}H_{15}N_5O$ requires 305.3). 1H -NMR (CD_3OD) δ : 2.33 (s, 3H, CH_3), 2.41 (s, 3H, CH_3), 6.74-6.56 (m, 3H, pyrimidinyl-H/Ph-H), 7.36 (d, 2H, $J = 8.5$ Hz, Ph-H), 8.25 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H).

3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile [34]

M.p. 195.6-198.9 °C. MS: $[M+H]^+ = 357.7$ ($C_{18}H_{14}F_3N_5$ requires 357.3). 1H -NMR ($CDCl_3$) δ : 2.33 (s, 3H, CH_3), 2.44 (s, 3H, CH_3), 6.75 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 7.20 (br. s, 1H, NH), 7.50 (d, 2H, $J = 8.8$ Hz, Ph-H), 7.71 (d, 2H, $J = 8.8$ Hz, Ph-H), 8.39 (d, 1H, $J = 5.1$ Hz, pyrimidinyl), 8.40 (br. s, 1H, NH).

4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
[35]

M.p. 178.3-181.2 °C. MS: $[M+H]^+ = 416.6$ ($C_{18}H_{14}IN_5$ requires 415.2). 1H -NMR ($CDCl_3$) δ : 2.39 (s, 3H, CH_3), 2.49 (s, 3H, CH_3), 6.76 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 7.10 (br. s, 1H, NH), 7.44 (d, 2H, $J = 8.8$ Hz, Ph-H), 7.61 (d, 2H, $J = 8.8$ Hz, Ph-H), 8.42 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 8.45 (br. s, 1H, NH).

4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [36]

M.p. 247-250 °C. MS: $[M+H]^+ = 305.8$ ($C_{17}H_{15}N_5O$ requires 305.3). 1H -NMR (DMSO- d_6) δ : 2.31 (s, 3H, CH_3), 2.41 (s, 3H, CH_3), 6.33 (m, 1H, Ph-H), 6.82 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 7.01 (t, 1H, $J = 8.1$ Hz, Ph-H), 7.11 (m, 1H, Ph-H), 7.33 (t, 1H, $J = 2.1$ Hz, Ph-H), 8.40 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.18 (s, 1H), 9.30 (s, 1H), 12.20 (br. s, 1H, NH).

3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile [37]

M.p. 233-237 °C. MS: $[M+H]^+ = 350.0$ ($C_{18}H_{16}N_6O_2$ requires 348.6). 1H -NMR (DMSO- d_6) δ : 2.31 (s, 3H, CH_3), 2.42 (s, 3H, CH_3), 2.44 (s, 3H, CH_3), 6.92 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.39 (d, 1H, $J = 8.5$ Hz, Ph-H), 7.87 (dd, 1H, $J = 8.1, 1.7$ Hz, Ph-H), 8.48 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 8.63 (d, 1H, $J = 1.7$ Hz, Ph-H), 9.87 (s, 1H, NH), 12.21 (br. s, 1H, NH).

4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [38]

M.p. 189.5-191.7 °C. MS: $[M+H]^+ = 431.5$ ($C_{18}H_{16}IN_5$ requires 429.6). 1H -NMR (DMSO- d_6) δ : 2.29 (s, 3H, CH_3), 2.32 (s, 3H, CH_3), 2.43 (s, 3H, CH_3), 6.85 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 7.20 (d, 1H, $J = 8.1$ Hz, Ph-H), 7.57 (m, 1H, Ph-H), 8.41-8.43 (m, 2H, Ph-H, pyrimidinyl-H), 9.48 (s, 1H, NH), 12.20 (br. s, 1H, NH).

4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [39]

M.p. 194.2-197.9 °C. MS: $[M+H]^+ = 338.0$ ($C_{18}H_{16}ClN_5$ requires 337.8). 1H -NMR (DMSO- d_6) δ : 2.27 (s, 3H, CH_3), 2.31 (s, 3H, CH_3), 2.41 (s, 3H, CH_3), 6.86 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.28 (d, 1H, $J = 8.5$ Hz, Ph-H), 7.61 (dd, 1H, $J = 8.8, 2.4$ Hz, Ph-H), 7.73 (d, 1H, $J = 2.7$ Hz, Ph-H), 8.43 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 9.51 (s, 1H, NH), 12.21 (br. s, 1H, NH).

4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [40]

M.p. 221-225 °C. MS: $[M+H]^+ = 320.9$ ($C_{18}H_{17}N_5O$ requires 319.4). 1H -NMR (DMSO- d_6) δ : 2.03 (s, 3H, CH_3), 2.29 (s, 3H, CH_3), 2.40 (s, 3H, CH_3), 6.78 (d, 1H, J = 5.1 Hz, pyrimidinyl-H), 6.89 (d, 1H, J = 8.1 Hz, Ph-H), 7.02 (dd, 1H, J = 8.3, 1.7 Hz, Ph-H), 7.29 (d, 1H, J = 0.7 Hz, Ph-H), 8.37 (d, 1H, J = 4.9 Hz, pyrimidinyl-H), 9.08 (s, 1H), 9.20 (s, 1H), 12.17 (br. s, 1H, NH).

4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [41]

M.p. 161.3-164.1 °C. MS: $[M+H]^+ = 321.6$ ($C_{18}H_{16}FN_5$ requires 321.4). 1H -NMR (DMSO- d_6) δ : 2.19 (s, 3H, CH_3), 2.30 (s, 3H, CH_3), 2.42 (s, 3H, CH_3), 6.82 (d, 1H, J = 5.1 Hz, pyrimidinyl-H), 7.03 (t, 1H, J = 9.3 Hz, Ph-H), 7.53 (m, 1H, Ph-H), 7.61 (dd, 1H, J = 7.1, 2.4 Hz, Ph-H), 8.39 (d, 1H, J = 5.1 Hz, pyrimidinyl-H), 9.36 (s, 1H, NH), 12.20 (br. s, 1H, NH).

4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [42]

M.p. 177.7-179.9 °C. MS: $[M+H]^+ = 322.5$ ($C_{18}H_{16}FN_5$ requires 321.3). 1H -NMR (DMSO- d_6) δ : 2.15 (s, 3H, CH_3), 2.30 (s, 3H, CH_3), 2.43 (s, 3H, CH_3), 6.86 (d, 1H, J = 5.1 Hz, pyrimidinyl-H), 7.13 (t, 1H, J = 9.0 Hz, Ph-H), 7.36 (dd, 1H, J = 8.1, 1.7 Hz, Ph-H), 7.75 (dd, 1H, J = 12.9, 1.5 Hz, Ph-H), 8.43 (d, 1H, J = 5.4 Hz, pyrimidinyl-H), 9.56 (s, 1H, NH), 12.21 (br. s, 1H, NH).


4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile [43]

M.p. 190.6-193.7 °C. MS: $[M+H]^+ = 334.7$ ($C_{19}H_{20}N_6$ requires 332.4). 1H -NMR ($CDCl_3$) δ : 2.36 (s, 3H, CH_3), 2.46 (s, 3H, CH_3), 2.94 (br. s, 6H, CH_3), 6.66 (d, 1H, J

= 5.6 Hz, *pyrimidinyl-H*), 6.79-6.80 (m, 2H, *Ph-H*), 7.05 (br. s, 1H, NH), 7.40-7.43 (m, 2H, *Ph-H*), 8.34 (d, 1H, $J = 5.1$ Hz, *pyrimidinyl-H*), 8.52 (br. s, 1H, NH).

Example 5

5 4-(3-Dimethylamino-acryloyl)-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide



1-(2,4-Dimethyl-1H-pyrrol-3-yl)-ethanone (1.1 g, 10 mmol) was partially dissolved in 2 M solution of ammonia in MeOH and H₂O₂ (10 mL of a 27 % w/w solution in H₂O) was added dropwise over a period of 40 min at such a rate as to maintain the internal
 10 temperature ≤ 30 °C. The mixture was stirred for 18 h at room temperature. The resulting suspended white solid was filtered and recrystallised from EtOAc to afford 4-acetyl-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide (1.06 g). An aliquot (720 mg, 4 mmol) was suspended in 1,1-bis-dimethylamino-3,3-dimethyl-butan-2-one (2mL, 9.6mmol) in a N₂-flushed flask and heated at 75 °C for 48 h. The crude mixture
 15 was cooled and purified by SiO₂ chromatography (EtOAc / MeOH gradient elution). The title compound (449 mg) was obtained as a buff solid. ¹H-NMR (DMSO-d₆) δ : 2.30 (s, 3H, CH₃), 2.46 (s, 3H, CH₃), 2.90 (br. s, 2H, NH). 3.09 (s, 3H, CH₃), 3.13 (s, 3H, CH₃), 5.23 (d, 1H, $J = 12.4$ Hz, CH), 7.38 (d, 1H, $J = 12.7$ Hz, CH), 10.97 (br. s, 1H, NH).

20

Example 6


4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide [44].

4-(3-Dimethylamino-acryloyl)-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide (100
 25 mg, 0.43 mmol), 4-fluorophenylguanidine nitrate (139 mg, 0.65 mmol) and K₂CO₃ (94 mg, 0.68 mmol) were partially dissolved in 2-methoxyethanol (5 mL) and heated at 120 °C for 18 h. The mixture was concentrated *in vacuo* and purified by SiO₂

chromatography (EtOAc / MeOH gradient elution). The crude product was triturated in iPr₂O to afford the title compound (31 mg) as a buff solid. M.p. 93.5-96.8 °C. MS: [M+H⁺] = 326.9 (C₁₇H₁₆FN₅O requires 325.3). ¹H-NMR (DMSO-d₆) δ: 2.36 (s, 3H, CH₃), 2.39 (s, 3H, CH₃), 6.79 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 6.92 (br. s, 2H, NH), 7.07 (t, 2H, *J* = 8.5 Hz, Ph-H), 7.76-7.78 (m, 2H, Ph-H), 8.36 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 9.41 (s, 1H, NH), 11.24 (br. s, 1H, NH).

Example 7

3-Dimethylamino-1-(3,5-dimethyl-1H-pyrrol-2-yl)-propenone



Pentane-2,4-dione (10.3 mL, 0.1 mol) was diluted with AcOH and cooled to 0 °C. NaNO₂ (6.9 g, 0.1 mol) was dissolved in H₂O (10 mL) and added dropwise, keeping the internal temperature ≤ 10 °C. The mixture was stirred for 1 h with cooling then 3 h at room temperature. Ethyl acetoacetate (14 mL, 0.11 mol) was dissolved in 1:1 AcOH / H₂O (100 mL) and Zn powder (7.19 g, 0.11 mol) was added. To this the oxime solution was added and the mixture was heated at 100 °C for 30 min then poured into H₂O (0.8 L). The aqueous mixture was extracted with EtOAc (3 × 500mL). The combined organic extracts were washed (brine), dried (MgSO₄), filtered, and evaporated *in vacuo*. The residue was purified by SiO₂ chromatography (heptane / EtOAc gradient elution) to afford 5-acetyl-2,4-dimethyl-1H-pyrrole-3-carboxylic acid ethyl ester (4.32 g). An aliquot (3.5 g, 16.7 mmol) was added to KOH (4.22 g, 75.3 mmol) dissolved in H₂O (5 mL) and the mixture was heated at 130 °C for 18 h. It was then cooled, diluted with H₂O (50 mL) and acidified with 2 M aq HCl. This mixture was extracted with EtOAc (3 × 50 mL). The combined organic extracts were washed (brine), dried (MgSO₄), filtered, and evaporated *in vacuo*. The residue was purified by SiO₂ chromatography (EtOAc) to afford 1-(3,5-dimethyl-1H-pyrrol-2-yl)-ethanone (1.05 g). An aliquot (536 mg, 3.9 mmol) was suspended in *N,N*-dimethylformamide dimethyl acetal (1.2 mL, 8.8 mmol) in a N₂-flushed flask and heated at 90 °C for 48 h.

The mixture was cooled and triturated in cold EtOAc. The resulting precipitate was filtered and washed with EtOAc to afford the title compound (444 mg) as a buff. ¹H-NMR (DMSO-d₆) δ: 1.41 (s, 3H, CH₃), 2.23 (s, 3H, CH₃), 2.23 (br. s, 6H, CH₃), 4.77 (d, 1H, *J* = 12.4 Hz, CH), 4.95 (s, 1H, pyrrolyl-H), 6.85 (d, 1H, *J* = 12.4 Hz, CH).

5

Example 8


[4-(3,5-Dimethyl-1H-pyrrol-2-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [45]

3-Dimethylamino-1-(3,5-dimethyl-1H-pyrrol-2-yl)-propenone (125 mg, 0.65 mmol), 4-fluorophenyl guanidine nitrate (211 mg, 0.98 mmol), and K₂CO₃ (149 mg, 1.08 mmol) were partially dissolved in 2-methoxyethanol and heated at 120 °C for 18 h. The mixture was concentrated *in vacuo* and purified by SiO₂ chromatography (5-g Isolute SI™ cartridge eluted with an heptane / EtOAc gradient). The crude product was triturated in iPr₂O to afford the title compound (158 mg) as a buff solid. M.p. 168.4-171.5 °C. MS: [M+H]⁺ = 283.9 (C₁₆H₁₅FN₄ requires 282.3). ¹H-NMR (CDCl₃) δ: 2.30 (s, 3H, CH₃), 2.35 (s, 3H, CH₃), 5.85 (s, 1H, pyrrolyl-H), 6.83 (d, 1H, *J* = 5.6 Hz, pyrimidinyl-H), 6.87 (br. s, 1H, NH), 7.05 (t, 2H, *J* = 8.5 Hz, Ph-H), 7.51-7.54 (m, 2H, Ph-H), 8.26 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 9.08 (br. s, 1H, NH).

15

Example 9

20 3-Dimethylamino-1-(1,2,4-trimethyl-1H-pyrrol-3-yl)-propenone



KOH (818 mg, 14.6 mmol) was partially dissolved in DMSO (15 mL) and stirred for 5 min. 1-(2,4-dimethyl-1H-pyrrol-3-yl)-ethanone (1 g, 7.3 mmol) was added in small portions and the mixture was stirred for 45 min. Iodomethane (0.54 mL, 8.75 mmol) was added dropwise at such a rate as to maintain the internal temperature ≤ 30 °C. The mixture was stirred for a further 45 min then poured into H₂O (50 mL) and extracted with Et₂O (3 × 60 mL). The combined organic extracts were washed (brine), dried

25


- (MgSO₄), filtered, and evaporated *in vacuo* to afford 1-(1,2,4-trimethyl-1*H*-pyrrol-3-yl)-ethanone (1.06 g) as a pink solid. This was suspended in 1,1-bis-dimethylamino-3,3-dimethyl-butan-2-one (3.6 mL, 17.5 mmol) in a N₂-flushed flask and heated at 70 °C for 36 h. The mixture was evaporated *in vacuo* and the residue triturated in EtOAc.
- 5 The titled compound (973 mg) was obtained as a reddish solid. ¹H-NMR (DMSO-d₆) δ: 1.30 (s, 3H, CH₃), 1.51 (s, 3H, CH₃), 2.20 (br. s, 6H, CH₃), 2.66 (s, 3H, CH₃), 4.57 (d, 1H, *J* = 12.5 Hz, CH), 5.53 (s, 1H, pyrrolyl-H), 6.72 (d, 1H, *J* = 12.7 Hz, CH).

Example 10

- 10 (4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-amine [46]
 3-Dimethylamino-1-(1,2,4-trimethyl-1*H*-pyrrol-3-yl)-propenone (150 mg, 0.73 mmol), 4-fluorophenyl guanidine nitrate (189 mg, 0.87 mmol) and K₂CO₃ (144 mg, 1.04 mmol) were partially dissolved in 2-methoxyethanol (3 mL) and heated at 110 °C 36 h. The mixture was concentrated *in vacuo* and purified by SiO₂ chromatography
- 15 (heptane / EtOAc gradient elution). The title compound (20 mg) was obtained as a brownish solid after recrystallisation from iPr₂O / heptane. M.p. 124.2-128.3 °C. MS: [M+H]⁺ = 298.4 (C₁₇H₁₇FN₄ requires 296.3). ¹H-NMR (CDCl₃) δ: 2.21 (s, 3H, CH₃), 2.42 (s, 3H, CH₃), 3.51 (s, 3H, CH₃), 6.40 (s, 1H, pyrrolyl-H), 6.76 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H), 7.01 (t, 2H, *J* = 8.8 Hz, Ph-H), 7.58-7.61 (m, 2H, Ph-H), 7.70 (br. s,
- 20 1H, NH), 8.28 (d, 1H, *J* = 5.6 Hz, pyrimidinyl-H).

Example 11

3-Dimethylamino-1-(2,4-dimethyl-5-nitro-1*H*-pyrrol-3-yl)-propenone



- 25 HNO₃ (0.28 mL of a 69 % w/v aq solution, 4.37 mmol) was added dropwise to Ac₂O (5 mL) at room temperature, keeping the internal temperature ≤ 25 °C. The nitrating

mixture was stirred at room temperature for 15 min before cooling to -40 °C. 1-(2,4-dimethyl-1*H*-pyrrol-3-yl)-ethanone (500 mg, 3.64 mmol) was dissolved in Ac₂O (6 mL) and added dropwise, keeping the internal temperature ≤ -30 °C. The mixture was stirred at -40 °C for 30 min then at -10 °C for a further 30 min. The mixture was poured into ice-water (50 mL) and was extracted with Et₂O (3 × 60 mL). The combined organic extracts were washed (brine), dried (Na₂SO₄), filtered, and evaporated *in vacuo* to give a dark brown solid. This was recrystallised from MeOH to afford 1-(2,4-dimethyl-5-nitro-1*H*-pyrrol-3-yl)-ethanone (158 mg). An aliquot (150 mg, 0.82 mmol) was suspended in 1,1-bis-dimethylamino-3,3-dimethyl-butan-2-one (0.42 mL, 2.02 mmol) in a N₂-flushed flask and was heated at 70 °C for 18 h. The mixture was triturated in EtOAc to afford the title compound (119 mg) as a brown solid. ¹H-NMR (DMSO-*d*₆) δ: 2.30 (s, 3H, CH₃), 2.40 (s, 3H, CH₃), 2.80 (br. s, 3H, CH₃), 3.07 (br. s, 3H, CH₃), 5.19 (d, 1H, *J* = 12.7 Hz, CH), 7.45 (d, 1H, *J* = 12.4 Hz, CH), 12.76 (br, 1H, NH).

15

Example 12

[4-(2,4-Dimethyl-5-nitro-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
[47]

3-Dimethylamino-1-(2,4-dimethyl-5-nitro-1*H*-pyrrol-3-yl)-propenone (110 mg, 0.46 mmol), 4-fluorophenyl guanidine nitrate (150 mg, 0.7 mmol), and K₂CO₃ (193 mg, 1.4 mmol) were partially dissolved in 2-methoxyethanol and heated at 120 °C for 18 h. The mixture was concentrated *in vacuo* and purified by SiO₂ chromatography (heptane / EtOAc gradient elution). The crude product was triturated in iPr₂O to afford the title compound (22 mg) as a pale orange solid. M.p. 166.3-170.1 °C. MS: [M+H]⁺ = 329.3 (C₁₆H₁₄FN₅O₂ requires 327.3). ¹H-NMR (DMSO-*d*₆) δ: 2.49 (s, 3H, CH₃), 2.59 (s, 3H, CH₃), 6.73 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 7.04 (t, 2H, *J* = 8.8 Hz, Ph-H), 7.07 (br. s, 1H, NH), 7.55-7.58 (m, 2H, Ph-H), 8.44 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 9.40 (br. s, 1H, NH).

The following compound was prepared in analogous manner:

N-[4-(2,4-Dimethyl-5-nitro-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-*N*',*N*'-dimethyl-benzene-1,4-diamine [48]

- 5 M.p. 265-268 °C. MS: $[M+H]^+ = 353.0$ ($C_{18}H_{20}N_6O_2$ requires 352.4). 1H -NMR (DMSO- d_6) δ : 2.39 (s, 3H, CH₃), 2.48 (br. s, 6H, CH₃), 2.82 (s, 3H, CH₃), 6.69 (d, 2H, $J = 9.0$ Hz, Ph-H), 6.74 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 7.50 (d, 2H, $J = 9.0$ Hz, Ph-H), 8.38 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 9.18 (s, 1H, NH), 13.00 (br. s, 1H, NH).

10

Example 13

[4-(5-Amino-2,4-dimethyl-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [49]

- [4-(2,4-Dimethyl-5-nitro-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine (45
15 mg, 0.14 mmol) was dissolved in EtOH (3 mL) and 10 % Pd(C) catalyst (10 mg) was added, followed by hydrazine hydrate (48 μ L of a 55 % w/w aq solution, 0.84 mmol). The mixture was heated at reflux for 18 h. The cooled mixture was filtered through a pad of Celite filter aid and the filtrate was evaporated *in vacuo*. The residue was purified by SiO₂ chromatography (20:1 EtOAc / 2 M ammonia in MeOH) to afford the
20 title compound (14 mg) as a yellow solid. M.p. 227-231 °C. MS: $[M+H]^+ = 397.8$ ($C_{16}H_{16}FN_5$ requires 297.3). 1H -NMR (CDCl₃) δ : 1.98 (s, 3H, CH₃), 2.37 (s, 3H, CH₃), 4.86 (br. s, 2H, NH₂), 6.65 (d, 1H, $J = 4.9$ Hz, pyrimidinyl-H), 7.03 (t, 2H, $J = 8.3$ Hz, Ph-H), 7.49 (br. s, 2H, NH), 7.58-7.61 (m, 2H, Ph-H), 8.53 (d, 1H, $J = 4.9$ Hz, pyrimidinyl-H).

25

Example 14

[4-(5-Bromo-2,4-dimethyl-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [50]

[4-(2,4-Dimethyl-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine (80 mg, 0.28 mmol) was dissolved in THF (4 mL) and cooled to -50 °C. *N*-Bromosuccinimide (55 mg, 0.31 mmol) was dissolved in THF (2 mL) and added dropwise, keeping the internal temperature \leq -40 °C. The mixture was stirred for 1h with cooling then
5 evaporated *in vacuo*. The residue was treated with H₂O (10 mL) and extracted with EtOAc (3 \times 10mL). The combined organic extractss were washed (brine), dried (MgSO₄), filtered, and evaporated *in vacuo*. The crude product was purified by SiO₂ chromatography heptane / EtOAc gradient elution) to afford the title compound (19 mg) as an orange solid after recrystallisation from iPr₂O. M.p. 181.4-183.3 °C. MS:
10 [M+H]⁺ = 362.9 (C₁₆H₁₄BrFN₄ requires 361.2). ¹H-NMR (CDCl₃) δ : 2.10 (s, 3H, CH₃), 2.36 (s, 3H, CH₃) 6.56 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 6.97 (t, 2H, *J* = 8.3Hz, Ph-H), 7.00 (br. s, 1H, NH), 7.79-7.52 (m, 2H, Ph-H), 8.85 (br. s, 1H, NH), 8.26 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H).

15 The following compound was prepared in analogous manner:

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
[51]

M.p. 198.1-203 °C. MS: [M+H]⁺ = 389.3 (C₁₆H₁₄BrN₅O₄ requires 361.2). ¹H-NMR
20 (CDCl₃) δ : 2.49 (s, 3H, CH₃), 2.59 (s, 3H, CH₃) 6.73 (d, 1H, *J* = 5.1 Hz, pyrimidinyl-H), 7.04 (t, 2H, *J* = 8.8 Hz, Ph-H), 7.57 (m, 2H, Ph-H), 7.90 (br. s, 1H, NH), 8.44 (d, 1H, *J* = 5.1Hz, pyrimidinyl-H), 9.40 (br. s, 1H, NH).

Example 15

25 *[4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine*
[52]

[4-(2,4-Dimethyl-1*H*-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine (80 mg, 0.28 mmol) was dissolved in THF (4 mL) and cooled to -60 °C. *N*-Chlorosuccinimide (41 mg, 0.31 mmol) was dissolved in THF (2 mL) and added dropwise, keeping the

internal temperature ≤ -50 °C. The mixture was stirred for 30min with cooling then evaporated *in vacuo*. The residue was treated with H₂O (10 mL) and extracted with EtOAc (3 \times 10 mL). The combined organic extracts were washed (brine), dried (MgSO₄), filtered, and evaporated *in vacuo*. The crude product was purified by SiO₂ chromatography (heptane / EtOAc gradient elution) to afford the title compound (37 mg) as an orange solid after recrystallisation from iPr₂O. M.p. 200-203 °C. MS: [M+H]⁺ = 317.7 (C₁₆H₁₄ClFN₄ requires 316.8). ¹H-NMR (CDCl₃) δ : 2.17 (s, 3H, CH₃), 2.42 (s, 3H, CH₃) 6.77 (d, 1H, *J* = 5.9 Hz, pyrimidinyl-H), 7.02-7.06 (m, 3H, Ph-h, NH), 7.54-7.56 (m, 2H, Ph-H), 7.95 (br. s, 1H, NH), 8.25 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H).

Example 16

[4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluorophenyl)-amine [53]

Diethylamine (40 μ L, 0.31 mmol) was diluted with methanol (0.5 mL) and formaldehyde (30 μ L of a 37 % w/w aq solution, 0.37 mmol) was added. [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluorophenyl)-amine (87 mg, 0.31 mmol) was added in small portions and the mixture was heated to reflux. After 1.5 h the mixture was diluted with H₂O (10 mL). The resulting precipitate was filtered and triturated in 2 M aq HCl. The mixture was filtered and the filtrate was washed with 2 M aq NaOH. The filtrate was extracted with EtOAc (3 \times 15 mL). The combined organic extracts were washed (brine), dried (MgSO₄), filtered, and evaporated *in vacuo*. The crude product was purified by SiO₂ chromatography (heptane / EtOAc gradient elution) to afford the title compound (36 mg) as an orange solid after recrystallisation from iPr₂O. M.p. 71.9-74.2 °C. MS: [M+H]⁺ = 367.7 (C₂₁H₂₆FN₅ requires 367.5). ¹H-NMR (CD₃OD) δ : 1.10 (t, 6H, *J* = 7.1 Hz, CH₃), 2.19 (s, 3H, CH₃), 2.41 (s, 3H, CH₃), 2.58 (t, 4H, *J* = 7.8 Hz, CH₂), 3.56 (s, 2H, CH₂), 6.78 (d, 1H, *J* = 5.6 Hz, pyrimidinyl-H), 7.00 (t, 2H, *J* = 8.5 Hz, Ph-H), 7.61-7.63 (m, 2H, Ph-H), 8.22 (d, 1H, *J* = 5.4 Hz, pyrimidinyl-H).

The following compounds was prepared in analogous manner:

5 *[4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [54]*

M.p. 88.4-91.6 °C. MS: $[M+H]^+ = 340.6$ ($C_{19}H_{22}FN_5$ requires 339.4). 1H -NMR ($CDCl_3$) δ : 2.18 (s, 3H, CH_3), 2.56 (s, 6H, CH_3), 2.42 (s, 3H, CH_3), 3.38 (s, 2H, CH_2), 6.75 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.00 (t, 2H, $J = 8.6$ Hz, Ph-H), 7.13 (br. s, 1H, NH), 7.56-7.59 (m, 2H, Ph-H), 8.31 (d, 1H, $J = 5.1$ Hz, pyrimidinyl-H), 8.55 (br. s, 1H, NH).

[4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine [55]

M.p. 94.7-97.6 °C. MS: $[M+H]^+ = 382.1$ ($C_{21}H_{24}FN_5O$ requires 381.5). 1H -NMR ($DMSO-d_6$) δ : 2.12 (s, 3H, CH_3), 2.33-2.35 (m, 7H, CH_3 , CH_2), 3.55 (m, 4H, CH_2), 4.03 (s, 2H, CH_2), 6.73 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 7.08 (t, 2H, $J = 9.0$ Hz, Ph-H), 7.74-7.78 (m, 2H, Ph-H), 8.28 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 9.27 (s, 1H, NH), 10.76 (s, 1H, NH).

20 *{4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine [56]*

M.p. 120.4-123.1 °C. MS: $[M+H]^+ = 396.4$ ($C_{22}H_{27}FN_5$ requires 394.5). 1H -NMR ($CDCl_3$) δ : 1.62 (br. s, 4H, CH_2), 2.10 (s, 3H, CH_3), 2.34 (s, 3H, CH_3), 2.37 (s, 3H, CH_3), 2.43 (br. s, 4H, CH_2), 3.42 (s, 2H, CH_2), 6.67 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 6.91-6.96 (m, 3H, Ph-H, NH), 7.50-7.52 (m, 2H, Ph-H), 8.24 (d, 1H, $J = 5.4$ Hz, pyrimidinyl-H), 8.30 (br. s, 1H, NH).

Example 17

Kinase specificity of selected compound

Selected compounds from the above examples were investigated for their kinase selectivity. A panel of protein kinases, including the CDKs relevant to the present invention, as well as a representative number of functionally unrelated kinases, were used.

Assays for CDK4/Cyclin D1, CDK2/Cyclin E, CDK1/Cyclin B kinase may be carried out by monitoring phosphorylation of GST-Rb in an appropriate system. Thus, GST-Rb phosphorylation, induced by CDK4/Cyclin D1, CDK2/Cyclin E or CDK1/Cyclin B is determined by incorporation of radio-labeled phosphate in GST-Rb(772-928) using radiolabelled ATP in 96-well format *in vitro* kinase assay. The phosphorylation reaction mixture (total volume 40 μ l) consisted of 50 mM HEPES pH 7.4, 20 mM MgCl₂, 5 mM EGTA, 2 mM DTT, 20 mM β -glycerophosphate, 2 mM NaF, 1 mM Na₃VO₄, Protease Inhibitors Cocktail (Sigma, see above), BSA 0.5mg/ml, 1 μ g purified enzyme complex, 10 μ l of GST-Rb-Sepharose beads, 100 μ M ATP, 0.2 μ Ci ³²P-ATP. The reaction is carried out for 30 min at 30°C at constant shaking. At the end of this period 100 μ l of 50 mM HEPES, pH 7.4 and 1 mM ATP is added to each well and the total volume transferred onto GFC filtered plate. The plate is washed 5 times with 200 μ l of 50 mM HEPES, pH 7.4 and 1 mM ATP. To each well were added 50 μ l scintillant liquid and the radioactivity of the samples is measured on Scintillation counter (Topcount, HP). The IC₅₀ values of different peptides were calculated using GraFit software.

Alternatively, CDK2/cyclin A kinase assays may be performed in 96-well plates using recombinant CDK2/cyclin A. Assay buffer consisted of 25 mM β -glycerophosphate, 20 mM MOPS, 5 mM EGTA, 1 mM DTT, 1mM NaVO₃, pH 7.4, into which is added 2 – 4 μ g of CDK2/cyclin A with substrate pRb(773-928). The reaction is initiated by addition of Mg/ATP mix (15mM MgCl₂, 100 μ M ATP with 30-50 kBq per well of

[γ - ^{32}P]-ATP) and mixtures incubated for 10 – 30 min, as required, at 30 °C. Reactions were stopped on ice, followed by filtration through p81 filterplates (Whatman Polyfiltronics, Kent, UK). After washing 3 times with 75 mM orthophosphoric acid, plates were dried, scintillant added and incorporated radioactivity measured in a scintillation counter (TopCount , Packard Instruments, Pangbourne, Berks, UK).

PKC α kinase activity may be measured by the incorporation of radio-labeled phosphate in Histone 3, as described. The reaction mixture (total volume 65 μl) consist of 50 mM Tris-HCl, 1 mM Calcium acetate, 3 mM DTT, 0.03 mg/ml Phosphatidylserine, 2.4 $\mu\text{g/ml}$ PMA, 0.04% NP40, 12 mM Mg/Cl, purified PKC α - 100 ng, Histone 3, 0.2mg/ml, 100 μM ATP, 0.2 μCi [γ - ^{32}P]-ATP. The reaction is carried over 15 min at 37°C in microplate shaker and is stopped by adding 10 μl 75 mM orthophosphoric acid and placing the plate on ice. 50 μl of the reaction mixture is transferred onto P81 filterplate and after washing off the free radioactive phosphate (3 times with 200 μl 75 mM orthophosphoric acid per well) 50 μl of scintillation liquid (Microscint 40) were added to each well and the radioactivity is measured on Scintillation counter (Topcount, HP).

For use in said assays CDK2 and/or PKC may be obtained from available sources or produced by recombinant methods as described. His-tagged CDK2/Cyclin E and CDK1/Cyclin B may be co-expressed and PKC α singularly expressed in Sf 9 insect cells infected with the appropriate baculovirus constructs. The cells are harvested two days after infection by low speed centrifugation and the proteins purified from the insect cell pellets by Metal-chelate chromatography. Briefly, the insect cell pellet is lysed in Buffer A (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.02% NP40 and 5 mM β -marcaptoethanol, 1 mM NaF, 1 mM Na_3VO_4 and Protease Inhibitors Cocktail (Sigma) containing AEBSF, pepstatin A, E 64, bestatin, leupeptin) by sonication. The soluble fraction is cleared by centrifugation and loaded onto Ni-NTA-Agarose (Quiagen). Non bound proteins were washed off with 300 mM NaCl, 5-15 mM

Imidazole in Buffer A and the bound proteins eluted with 250 mM Imidazole in Buffer A. The purified proteins are extensively dialyzed against Storage buffer (20 mM HEPES pH 7.4, 50 mM NaCl, 2 mM DTT, 1 mM EDTA, 1 mM EGTA, 0.02% NP40, 10% v/v Glycerol) aliquoted and stored at -70°C. PKC- α - 6 x His may be
 5 purified the same way but using different buffers- 50 mM NaH₂PO₄, pH 8.0 and 0.05% Triton X-100 instead of Tris and NP40 respectively.

The results in the Table 1 below show that the compounds in question exhibit a high degree of selectivity for inhibition of CDKs.

10

Table 1

Compound	CDK2/cyclin E (IC ₅₀ , μ M)	Compound	CDK2/cyclin E (IC ₅₀ , μ M)
1	1.0 \pm 0.7	38	0.86
2	0.04	39	1.02
3	0.5	40	0.15
4	1.3 \pm 0.4	41	0.05
5	1.7	42	0.76
6	1.3	43	0.54
7	2.9	44	0.22
8	1.4	46	1.79
12	0.90	47	0.06
16	1.52	48	0.96
17	1.73	50	0.55
31	0.03	51	0.37
32	0.09	52	0.73
33	0.09	53	0.87
34	0.18	54	0.14
35	0.55	55	0.87

36	0.04	56	1.51
37	0.39		

Example 18

Anti-proliferative effect of selected compounds

Selected compounds from the above examples were subjected to a standard cellular proliferation assay using a range of different human tumour cell lines. Standard 72-h MTT (thiazolyl blue; 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assays were performed (Haselsberger, K.; Peterson, D. C.; Thomas, D. G.; Darling, J. L. Anti Cancer Drugs 1996, 7, 331-8; Loveland, B. E.; Johns, T. G.; Mackay, I. R.; Vaillant, F.; Wang, Z. X.; Hertzog, P. J. Biochemistry International 1992, 27, 501-10).

Human tumour cell lines were obtained from the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, USA).


The results in Table 2 below illustrate the anti-proliferative effect of compounds described in this application.

Table 2

	Cytotoxicity (IC ₅₀ , μ M)							
	Human tumour cell line					Non-transformed cell line		
Compound	A549	HeLa	HT29	MCF7	Saos-2	Hs27	IMR90	W138
2	3.30		3.30		6.60			
12	15.00		3.10		17.00			
16	4.39							
31	0.40	0.26	0.31	0.26	0.74	18.9	14.1	9.7
32	2.47	1.38	2.41	1.93	2.89	15.1	20.8	9.6
33	0.88	0.76	1.83	1.03	0.89	4.2	14.1	9.9
34	3.64		0.95		3.40			
35	0.51	0.09	0.11	0.45	0.75	35.1	37.3	21.3
36	1.12		2.56		0.90			
37	0.47		0.28		1.04			
38	3.60		0.72		3.12			
39	4.37		3.27		5.88			
40	0.86		1.47		1.13			
41	2.95		2.11		5.80			
42	1.49		0.28		2.23			
43	1.41		0.17		1.50			
44	1.36		1.92		2.61			
47	3.13		3.11		3.78			
48	3.05		0.61		2.65			
51	7.56		2.92		12.40			
53	0.97		1.31		1.61			
54	2.76		3.20		0.70			
56	13.80		5.09		14.15			

CLAIMS

1. A compound of general formula I:



I

wherein:

one of X^1 and X^2 is NR^{10} and the other of X^1 and X^2 is CR^9 ;

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, or CH=CH;

R^1 , R^2 , R^3 , R^9 and R^{10} are independently H, alkyl, aryl, aralkyl, heterocycle, halogeno, NO₂, CN, OH, alkoxy, aryloxy, $(R''')_nNH_2$, $(R''')_nNH-R'$, $(R''')_nN-(R')(R'')$, NH-aryl, N-(aryl)₂, COOH, COO-R', COO-aryl, CONH₂, CONH-R', CON-(R')(R''), CONH-aryl, CON-(aryl)₂, SO₃H, SO₂NH₂, CF₃, CO-R', or CO-aryl, wherein alkyl, aryl, aralkyl and heterocycle groups may be further substituted with one or more groups selected from halogeno, NO₂, CN, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;

R^4 , R^5 , R^6 , R^7 , and R^8 are independently from each other H, substituted or unsubstituted lower alkyl, halogeno, NO₂, CN, OH, substituted or unsubstituted alkoxy, NH₂, NH-R', N-(R')(R''), COOH, COO-R', CONH₂, CONH-R', CON-(R')(R''), SO₃H, SO₂NH₂, or CF₃;

wherein R', R'' and R''' are each independently alkyl groups that may be the same or different and n is 0 or 1;

with the proviso that when R¹ and R² are H, X¹ is NH, X² is CH, and R³ is H, the phenyl group is not

- 5 unsubstituted,
- 4-ethyl,
- 3-methyl,
- 3-(1,1,2,2- tetrafluoroethoxy),
- 3,4,5-trimethoxy,
- 10 when the other groups R⁴-R⁸ are H;
- and pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein;

- 15 - X¹ and X² are CR⁹ and NH respectively;
- R¹, R², R³ and R⁹ are each independently selected from H, alkyl, aryl, aralkyl, heterocycle, halogeno, NO₂, CN, OH, alkoxy, aryloxy, (R''')_nNH₂, (R''')_nNH-R', (R''')_nN-(R')(R''), COOH, COO-R', CONH₂, CONH-R', CON-(R')(R''), SO₃H,
- 20 SO₂NH₂, CF₃, and CO-R' wherein alkyl, aryl and aralkyl groups may be further substituted with one or more groups selected from halogeno, NO₂, CN, OH, O-methyl, NH₂, COOH, CONH₂ and CF₃;
- Z is selected from NH, NHSO₂ and NHCH₂;
- 25 - R⁴-R⁸ are each independently selected from H, halogeno, nitro, amino, aminoalkyl, hydroxy, alkoxy, carbamoyl, sulfamyl, N(R')(R''), C₁₋₄ alkyl and substituted C₁₋₄ alkyl.

3. A compound according to any preceding claim, wherein Z is NH and R³ is H.
4. A compound according to claim 3, wherein R¹, R² and R⁹ are each independently H, halogeno, CN, NO₂, CO(NH₂), (R''')NH(R')(R'') a C₁₋₄ alkyl group or a
5 heterocyclic group.
5. A compound according to claim 4, wherein when R¹ is halogeno, it is selected from chloro or bromo; when R¹ is alkylamino, it is diethylaminomethyl or dimethylaminomethyl; when R¹ is a heterocyclic group it is morpholin-4-ylmethyl or
10 4-methyl-piperazin-1-ylmethyl.
6. A compound according to any of claims 1-5, wherein R¹ is H or CN, and R² and R⁹ are both methyl.
- 15 7. A compound according to claim 7, wherein R¹ is H.
8. A compound according to claim 7, wherein R¹ is CN.
9. A compound according to any preceding claim, wherein;
20 R⁴, R⁵, R⁶, R⁷, and R⁸ are independently from each other H, unsubstituted lower alkyl, halogeno, NO₂, CN, OH, N-(R')(R''), or CF₃;
wherein R' R'' and R''' are each independently alkyl groups that may be the same or different and n is 0 or 1;
- 25 10. A compound according to claim 9, wherein R⁴ to R⁸ are selected independently from H, F, NH₂, NO₂, OH, Cl, Br, I, CN, CH₂OH, CF₃ and dimethylamino.
11. A compound according to claim 9 or 10, wherein R⁴ and R⁸ are both hydrogen.

12. A compound according to any preceding claim selected from 2-[N-(phenyl)]-4-(2,4-dimethylpyrrol-3-yl)pyrimidineamines in which the phenyl group is 2-, 3-, 4- or 5-substituted by at least one of F, NH₂, NO₂, OH, Cl, Br, I, CN, CH₂OH, CF₃ or OMe.

5

13. A compound according to claim 12, wherein the phenyl group is mono-substituted by F, NH₂, NO₂, OH, Cl, Br, I, CH₂OH, CN, CF₃ or OMe at any of the 2,3, 4 or 5-positions, or di-substituted by 2,4-difluoro, 3,5-difluoro, 3,4-difluoro, 2,4-dichloro, 3,5-dichloro, 3,4-dichloro or 4-chloro-3-trifluoromethyl.

10

14. A compound according to any of claims 1-11 selected from 2-[N-(phenyl)]-4-(3,5-dimethyl-1H-pyrrole-2-carbonitrile)pyrimidineamines in which the phenyl group is 2-, 3- or 4-substituted by at least one of F, NH(CH₃)₂, NO₂, OH, Cl, Br, I or CF₃.

15 15. A compound according to claim 14, wherein the phenyl group is mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at any of the 3 or 4-positions, or di-substituted by 4-methyl-3-nitro, 3-iodo-4-methyl, 4-chloro-3-methyl, 3-hydroxy-4-methyl, 4-fluoro-3-methyl or 4-methyl-3-fluoro.

20 16 A compound according to any of claims 1-5 selected from 2-[N-(phenyl)]-4-(2,4-dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position.

25 17. A compound according to claim 16, wherein the phenyl group is substituted by a fluoro or NH(CH₃)₂ group.

18. A compound according to any of claims 1-5 selected from 2-[N-(phenyl)]-4-(2,4-dimethyl-5-halogeno-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 3 or 4-positions.

19. A compound according to claim 18, wherein the phenyl group is substituted by a 4-fluoro or 3-nitro group, the halogeno group being chloro or bromo.
- 5 20. A compound according to any of claims 1-5 selected from 2-[N-(phenyl)]-4-(2,4-dimethyl-5-dialkylaminoalkyl-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group is mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position.
21. A compound according to claim 20, wherein the phenyl group is substituted by
10 fluoro, the dialkylaminoalkyl group preferably being diethylaminomethyl or dimethylaminomethyl.
22. A compound according to any of claims 1-5 selected from 2-[N-(phenyl)]-4-(2,4-dimethyl-5-(heterocycle)-1H-pyrrol-3-yl)-pyrimidinamines wherein the phenyl group
15 is preferably mono-substituted by F, NH(CH₃)₂, NO₂, OH, I or CF₃ at the 4-position.
23. A compound according to claim 22, wherein the phenyl group is substituted by fluoro, the heterocycle group being 5-morpholin-4-ylmethyl or 4-methyl-piperazin-1-ylmethyl.
- 20 24. A compound according to claim 1 selected from;
- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine
25 (3,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
(4-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
(3,5-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
4-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
3-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol

- (2,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (2,4-Dichloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (4-Chloro-3-trifluoromethyl-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
- 5 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-trifluoromethyl-phenyl)-amine
 (3-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 N-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine
- 10 (3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-fluoro-4-iodo-phenyl)-amine
 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
- 15 carbonitrile
 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
- 20 carbonitrile
 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 25 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 5 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- 4-(3,5-Dimethyl-1H-pyrrol-2-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 10 (4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine
- [4-(5-Amino-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 15 [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 20 [4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- {4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine.
- 25

25. A compound according to claim 24 selected from;

- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine
 (3,4-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (4-Chloro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 (3,5-Difluoro-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 5 4-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
 3-[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-ylamino]-phenol
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
 (3-Chloro-4-iodo-phenyl)-[4-(2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-fluoro-4-iodo-phenyl)-amine
 10 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-
 15 carbonitrile
 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-
 20 carbonitrile
 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 25 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile
 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-
 carbonitrile

- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 5 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- (4-Fluoro-phenyl)-[4-(1,2,4-trimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-amine
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-
- 10 1,4-diamine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-
- 15 phenyl)-amine
- [4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine, and
- 20 {4-[2,4-Dimethyl-5-(4-methyl-piperazin-1-ylmethyl)-1H-pyrrol-3-yl]-pyrimidin-2-yl}-(4-fluoro-phenyl)-amine.

26. A compound according to claim 25 selected from;

- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 25 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine
- [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-trifluoromethyl-phenyl)-amine
- 3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 5 4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile
- 10 4-[2-(3-Iodo-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Chloro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 15 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(3-Fluoro-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 20 4-[2-(4-Dimethylamino-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile
- 4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide
- [4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- 25 N-[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-N',N'-dimethyl-benzene-1,4-diamine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine
- [4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine
- [4-(5-Chloro-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Diethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine, and

- 5 [4-(2,4-Dimethyl-5-morpholin-4-ylmethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine.

27. A compound according to claim 26 selected from;

[4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine

- 10 [4-(2,4-Dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-iodo-phenyl)-amine

3,5-Dimethyl-4-[2-(3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(4-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 15 3,5-Dimethyl-4-[2-(4-trifluoromethyl-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(4-Iodo-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(3-Hydroxy-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 20 3,5-Dimethyl-4-[2-(4-methyl-3-nitro-phenylamino)-pyrimidin-4-yl]-1H-pyrrole-2-carbonitrile

4-[2-(3-Hydroxy-4-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

- 25 4-[2-(4-Fluoro-3-methyl-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carbonitrile

4-[2-(4-Fluoro-phenylamino)-pyrimidin-4-yl]-3,5-dimethyl-1H-pyrrole-2-carboxylic acid amide

[4-(2,4-Dimethyl-5-nitro-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine

[4-(5-Bromo-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(3-nitro-phenyl)-amine,
and

[4-(5-Dimethylaminomethyl-2,4-dimethyl-1H-pyrrol-3-yl)-pyrimidin-2-yl]-(4-fluoro-phenyl)-amine.

5

28. A compound according to claim 1, wherein;

- X^1 and X^2 are NH and CR^9 respectively;

- R^1 , R^2 , R^3 and R^9 are each independently selected from H, alkyl, aryl, aralkyl;
10 heterocycle, halogeno, NO_2 , CN, OH, alkoxy, aryloxy, $(R''')_nNH_2$, $(R''')_nNH-R'$,
 $(R''')_nN-(R')(R'')$, $COOH$, $COO-R'$, $CONH_2$, $CONH-R'$, $CON-(R')(R'')$, SO_3H ,
 SO_2NH_2 , CF_3 , and $CO-R'$ wherein alkyl, aryl and aralkyl groups may be further
substituted with one or more groups selected from halogeno, NO_2 , CN, OH, O-methyl,
 NH_2 , $COOH$, $CONH_2$ and CF_3 ;

15

- Z is selected from NH, $NHSO_2$ and $NHCH_2$;

- R^4 , R^5 and R^8 are each independently selected from H, halogeno, nitro, amino,
aminoalkyl, hydroxy, alkoxy, carbamoyl, sulfamyl, $N(R')(R'')$, C_{1-4} alkyl and
20 substituted C_{1-4} alkyl;

- R_6 is selected from H, halogeno, nitro, amino, aminoalkyl, hydroxy, alkoxy,
carbamoyl, sulfamyl, $N(R')(R'')$, methyl, propyl, butyl and substituted C_{1-4} alkyl;

- R_7 is selected from H, halogeno, nitro, amino, aminoalkyl, hydroxy, carbamoyl,
sulfamyl, $N(R')(R'')$ C_{2-4} alkyl and substituted C_{1-4} alkyl

25

29. Pharmaceutical compositions comprising a compound as defined in any of claims
1 to 28 or a pharmaceutically acceptable salt thereof together with a pharmaceutically
acceptable excipient.

30. Use of a compound as defined in any of claims 1 to 28 or a pharmaceutically acceptable salt thereof in the treatment of a proliferative disorder.


5 31. Use according to claim 30, wherein the proliferative disorder is cancer or leukaemia.

32. Use according to claim 30 or 31, wherein said compound is administered in an amount sufficient to inhibit at least one CDK enzyme.

10

33. Use according to claim 32, wherein the CDK enzyme is CDK2 and/or CDK4.

34. Use of a compound of formula



15

wherein:

20 one of X^1 and X^2 is NR^{10} and the other of X^1 and X^2 is CR^9 ;

Z is NH, NHCO, NHSO₂, NHCH₂, CH₂, CH₂CH₂, or CH=CH;

R^1 , R^2 , R^3 , R^9 and R^{10} are independently H, alkyl, aryl, aralkyl, heterocycle, halogeno,
 25 NO₂, CN, OH, alkoxy, aryloxy, $(R''')_nNH_2$, $(R''')_nNH-R'$, $(R''')_nN-(R')(R'')$, NH-

aryl, N-(aryl)₂, COOH, COO-R', COO-aryl, CONH₂, CONH-R', CON-(R')(R''),
 CONH-aryl, CON-(aryl)₂, SO₃H, SO₂NH₂, CF₃, CO-R', or CO-aryl, wherein alkyl,
 aryl, aralkyl and heterocycle groups may be further substituted with one or more
 groups selected from halogeno, NO₂, CN, OH, O-methyl, NH₂, COOH, CONH₂ and
 5 CF₃;

R⁴, R⁵, R⁶, R⁷, and R⁸ are independently from each other H, substituted or
 unsubstituted lower alkyl, halogeno, NO₂, CN, OH, substituted or unsubstituted
 alkoxy, NH₂, NH-R', N-(R')(R''), COOH, COO-R', CONH₂, CONH-R', CON-
 10 (R')(R''), SO₃H, SO₂NH₂, or CF₃;

wherein R' R'' and R''' are each independently alkyl groups that may be the same or
 different and n is 0 or 1;
 with the proviso that when R¹ and R² are H, X¹ is NH, X² is CH, and R³ is H, the
 15 phenyl group is not

3-(1,1,2,2- tetrafluoroethoxy), or

3,4,5-trimethoxy,

when the other groups R⁴-R⁸ are H;

and pharmaceutically acceptable salts thereof;












20 in the manufacture of a medicament for use in the treatment of a proliferative disease.












35. Use according to claim 34, wherein the compound is as defined in any of claims 2
 to 13.











25

Figure 1

No.	Structure
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	

12	
13	
14	
15	
16	
17	
31	
32	
33	
34	
35	

36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	

47	
48	
49	
50	
51	
52	
53	
54	
55	
56	

INTERNATIONAL SEARCH REPORT

In national Application No

EP/GB 02/01445

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D403/04 A61K31/506 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 233 461 A (AMERICAN CYANAMID CO) 26 August 1987 (1987-08-26) cited in the application page 8, line 9; claim 1 page 11, line 15 page 26, line 19-26 ---	1, 9-11, 29
X	WO 97 19065 A (CELLTECH THERAPEUTICS LTD ;DAVIS PETER DAVID (GB); MOFFAT DAVID FE) 29 May 1997 (1997-05-29) claim 1; example 14 ---	1-35
X	WO 95 09852 A (CIBA GEIGY AG) 13 April 1995 (1995-04-13) claims 1,6 ---	1-35
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

13 June 2002

Date of mailing of the international search report

21/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Grassi, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/01445

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 588 762 A (CIBA GEIGY AG) 23 March 1994 (1994-03-23) claim 1 ---	1-35
A	WO 00 39101 A (BREAULT GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06) the whole document -----	1-35

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/01445

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0233461	A	26-08-1987	AT 135699 T	15-04-1996
			AU 621461 B2	12-03-1992
			AU 5057890 A	26-07-1990
			AU 591223 B2	30-11-1989
			AU 6751887 A	16-07-1987
			BR 1100989 A3	09-11-1999
			CA 1320201 A1	13-07-1993
			DE 3751742 D1	25-04-1996
			DE 3751742 T2	21-11-1996
			DK 15187 A	14-07-1987
			EP 0233461 A2	26-08-1987
			ES 2087056 T3	16-07-1996
			FI 870113 A ,B,	14-07-1987
			GR 3019455 T3	30-06-1996
			HU 43582 A2	30-11-1987
			IE 74202 B1	16-07-1997
			JP 2043824 C	09-04-1996
			JP 7080857 B	30-08-1995
			JP 62223177 A	01-10-1987
			KR 9004693 B1	02-07-1990
			NZ 218924 A	26-04-1990
			PH 25056 A	28-01-1991
			SG 47583 A1	17-04-1998
			US 4876252 A	24-10-1989
			US 4788195 A	29-11-1988
			ZA 8700219 A	26-08-1987
WO 9719065	A	29-05-1997	AU 7631496 A	11-06-1997
			EP 0862560 A1	09-09-1998
			WO 9719065 A1	29-05-1997
			US 6235746 B1	22-05-2001
			US 5958935 A	28-09-1999
WO 9509852	A	13-04-1995	AU 693804 B2	09-07-1998
			AU 7697594 A	01-05-1995
			CA 2148477 A1	13-04-1995
			WO 9509852 A1	13-04-1995
			EP 0672040 A1	20-09-1995
			JP 8504834 T	28-05-1996
			US 5521184 A	28-05-1996
			US 5543520 A	06-08-1996
EP 0588762	A	23-03-1994	EP 0588762 A1	23-03-1994
			JP 6184116 A	05-07-1994
			US 5516775 A	14-05-1996
WO 0039101	A	06-07-2000	AU 1874300 A	31-07-2000
			BR 9916590 A	23-10-2001
			CN 1335838 T	13-02-2002
			EP 1140860 A1	10-10-2001
			WO 0039101 A1	06-07-2000
			NO 20013038 A	22-08-2001

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 2002/079193 A1

(51) International Patent Classification⁷: **C07D 403/04**,
A61K 31/506, A61P 35/00

(21) International Application Number:
PCT/GB2002/001445

(22) International Filing Date: 26 March 2002 (26.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0107901.1 29 March 2001 (29.03.2001) GB

(71) Applicant (for all designated States except US): **CYCLACEL LIMITED** [GB/GB]; 12 St. James's Square, London SW1Y 4RB (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FISCHER, Peter, Martin** [CH/GB]; Denley Lodge, 1 Arbirlot Road, Arbroath, Angus DD11 2EN (GB). **WANG, Shudong** [AU/GB]; Burnside Mill, Forfar, Angus, Scotland DD8 2RZ (GB). **WOOD, Gavin** [GB/GB]; 31 Millbank, Cupar, Fife, Scotland KY15 5DP (GB).

(74) Agents: **NACHSHEN, Neil, Jacob et al.**; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

(48) Date of publication of this corrected version:

17 March 2005

(15) Information about Correction:

see PCT Gazette No. 11/2005 of 17 March 2005, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF CYCLIN DEPENDENT KINASES AS ANTI-CANCER AGENT

(57) Abstract: The present invention relates to 2-substituted 4-heteroaryl-pyrimidines, their preparation, pharmaceutical compositions containing them and their use as inhibitors of cyclin-dependent kinases (CDKs) and hence their use in the treatment of proliferation disorders such as cancer, leukaemia, psoriasis and the like.

WO 2002/079193 A1